



PROGRAMME REGIONAL DE PRODUCTION INTEGREE DU COTON EN AFRIQUE

Bénin, Burkina Faso, Cameroun,
Côte d'Ivoire, Mali, Sénégal,
Tchad et Togo

Tél : (+226) 76 59 55 01/ 70 07 73 51

E-mail : prpica@yahoo.fr

Site Web : www.prpica.org

INFOS PR-PICA

PR-PICA NEWSLETTER

N°64

July 2026

CONTENTS

2026/2027 COTTON SEASON :

- ✚ Rainfall data: July 2026 **P. 2**
- ✚ Areas sown on July 2026 **P. 3**
- ✚ Pest situation as of July 31, 2026 **P. 4**
- ✚ Phytosanitary treatment program **P. 5-7**
- ✚ List of effective insecticides against jassids **P. 8**

**Newsletter published by
the Executive Secretariat of
PR-PICA.**

- **Chairman of PR-PICA :**
M. Tete AWOKOU
- **Vice-Chairman :**
Dr Abdou TRAORE
- **Rapporteur :**
M. NADAMA
- **Executive Secretary :**
M. Félix SAWADOGO



Rainfall levels recorded in July 2026 were average across all PR-PICA countries, with pockets of drought observed in several countries.

Crop area forecasts vary from one country to another. Several countries, such as Côte d'Ivoire, Chad and Togo, have exceeded their crop area forecasts. Rainfall shortfalls are largely to blame for the other countries failing to meet their forecasts.

Pest pressure as at 31 July was generally low for carpophagous pests, although there were localised moderate infestations of *H. armigera* and *Earias* in some countries. As for sucking pests, low levels of infestation were also noted, except in Chad, where the new species of jassid, *Amrasca biguttula*, was present.

The phytosanitary treatment products used this season often vary from one country to another within the same treatment window, but are in line with the PR-PICA recommendations.

RAINFALL FOR THE MONTH OF JULY 2026

Agro-ecological zones	Dekad	Bénin		Burkina Faso		Cameroun		Côte d'Ivoire		Mali		Sénégal		Tchad		Togo	
		nber days	height (mm)	nber days	height (mm)	nber days	height (mm)	nber days	height (mm)	nber days	height (mm)	nber days	height (mm)	nber days	height (mm)	nber days	height (mm)
Dry zone/ North	Dekad 1	5	29	3	75	4	53	2	31	4	131	1	14	1	23	3	57
	Dekad 2	5	36	2	46	6	186	5	47	3	22	3	60	3	73	3	75
	Dekad 3	4	60	4	92	4	90	5	84	2	16	2	21	3	61	3	70
	Total	14	124	9	212	14	329	12	162	9	169	6	95	7	157	10	202
Median zone/ Center	Dekad 1	5	36			3	52	2	14	1	4	1	16	2	33	3	64
	Dekad 2	6	67			5	64	5	95	4	40	5	97	3	48	4	108
	Dekad 3	5	26			5	113	6	116	3	54	4	78	3	64	4	70
	Total	16	129			13	229	13	225	8	98	9	191	8	145	11	243
Humid zone South	Dekad 1	3	22	2	43	1	12	2	39	3	123	2	22	2	59	1	15
	Dekad 2	5	43	3	59	5	170	1	29	4	50	5	103	3	58	3	62
	Dekad 3	1	7	3	60	4	79	2	35	3	31	4	119	3	71	2	34
	Total	9	72	8	162	10	261	5	103	10	204	11	244	8	189	6	112
AVERAGE JULY 2026		13	108	9	187	12	273	10	163	9	157	9	177	8	163	9	186
RAINFALL DISTRIBUTION		XX		XX		XXX		XX		XX		XX		XX		XXX	

X= Low distribution. XX : Average distribution. XXX : Good distribution


Rainfall levels were average across all countries, with totals ranging from 108 mm in Benin to 273 mm in Cameroon.

As for the distribution of rainfall, it was average to good, ranging from 8 days in Chad to 13 days in Benin.

Several countries experienced rainfall-related difficulties. This was the case in Benin, Burkina Faso, Cameroon and Mali, where pockets of drought were recorded.

This situation hampered the achievement of the season's sowing forecasts in these countries.

Compared with rainfall in July of the 2025/2026 season, a decrease in rainfall was noted for July 2026 in Benin (-63 mm), Burkina Faso (-4 mm), Côte d'Ivoire (-6 mm), Mali (-96 mm), Senegal (-6 mm) and Chad (-50 mm). However, an increase was recorded in Cameroon (+16 mm) and Togo (+99 mm).



Rainfall in July 2026 compared to the last two seasons

AREA SOWN, 2026/2027 SEASON

	Bénin	Burkina Faso	Cameroun	Côte d'Ivoire	Mali	Sénégal	Tchad	Togo
FORECAST 2026/2027 SEASON (ha)	600 000	532 046	209 676	351 000	630 000	30 000	212 908	105 000
Total area sown as of July 31, 2026 (ha)	519 596	462 839	201 507	351 616	488 631	29 888	214 008	115 980
Implementation rate as of July 31, 2026 (%)	86,6	86,9	96,1	100,2	77,6	99,6	101	110,5
Percentage of sowing for the month of July 2026 /Total area sown (%)	14	40	6	12	23	49	14	55
Evolution of area compared to the previous season (%)	0,7	9,6	1,2	21,2	-22	38,2	42,6	59,1

NB: These areas as of July 31, 2026 are not definitive and may be subject to readjustment..


As at 31 July 2026, the areas sown compared with the season's forecasts vary from one country to another, with achievement rates ranging from 77.6% in Mali to 110.5% in Togo. Côte d'Ivoire, Chad and Togo are the countries that have met or exceeded their season's forecasts. It should be noted that Burkina Faso has revised its crop area forecasts for the season upwards.

Several difficulties have led to the failure to meet crop area forecasts in the other countries.

- In Benin, pockets of drought disrupted sowing, causing some delays in all production zones;
- in Burkina Faso, an uneven spatial and temporal distribution of rainfall was observed, with two successive long periods of drought lasting between 10 and 17 days in the central region;
- in Mali, in addition to the rainfall deficit and delays in the delivery of fertiliser, poor seed quality was noted in the Fana and Bougouni regions.

The percentage of areas sown in July (late sowing) ranges from low to moderate depending on the country.

Compared with the previous 2025/2026 season, as at 31 July of this season, an increase in the area under cultivation was noted in all countries except Mali (-22 %, representing a decrease of 137,650 ha).



**Areas sown compared: 2024/2025, 2025/2026
and 2026/2027 seasons**

AVERAGE PEST SITUATION AT 31 JULY 2026

Pests	Infestation level by country								Observations
	Bénin	Burkina Faso	Cameroun	Côte d'Ivoire	Mali	Sénégal	Tchad	Togo	
<i>H. armigera</i>	X	X	XX	X	X	X	X	X	Average infestations in Cameroun
<i>Earias spp</i>	X		XX	X	X	X	X	X	Average infestations in Cameroun
<i>D. watersi</i>	X		X	X	X	X	X	X	Low infestations
<i>T. leucotreta</i>	X			X			X		Low infestations
<i>P. gossypiella</i>	X			X			X		Low infestations
<i>Bemisia tabaci</i>	X		X	X	X	X	X	X	Low infestations
Jassides (Amrasca biguttula)	X	X	X	X	X	X	XX	X	Average infestations in Tchad
Jassides (Autres)	X	X	X	X	X	X	X	X	Low infestations
<i>Dysdercus spp</i>				X	X	X	X	X	Low infestations
<i>A. gossypii</i>	X		X	X	X	X	X	X	Low infestations
<i>P. latus</i>	X	X	X	X	X	X	X	X	Low infestations
<i>H. derogata</i>	X		X	X	X	X	X	X	Low infestations
<i>A. flava</i>	X	X		X	X	X	X	X	Low infestations
<i>S. littoralis</i>	X		X	X	X	X	X	X	Low infestations
Virescence florale				X			X		Low infestations
Fusariose	X			X		X			Low infestations
Bactériose	X								Low infestations
<i>S. frugiperda (sur coton)</i>	X			X					Low infestations

X = Low infestation; XX = Medium infestation; XXX = High infestation

Pest pressure was low in July for carpophagous insects in most countries. However, in Cameroon, moderate infestations of *H. armigera* and *Earias spp* were noted on early-season seedlings.

As for sucking pests, low levels of infestation were also observed across all countries. The jassid, *Amrasca biguttula*, unlike in previous years, is showing low levels of infestation at the start of the season, except in Chad where moderate infestations have been observed. The use of jassid-control products recommended by the PR-PICA has made it possible to control infestations of this pest overall.

In Côte d'Ivoire, moderate levels of floral virescence have been noted in some COIC areas (the north-central part of the cotton-growing region). The same applies to fusarium wilt in CIDT areas (the southern part of the cotton-growing region).

For effective management of these pests, particularly jassids, we recommend:

-  **increased monitoring in cotton fields;**
-  **strict adherence to the established crop protection treatment schedule;**
-  **a smooth transition between crop protection treatment windows;**
-  **the use of recommended and appropriate insecticides (taking particular account of the modes of action of the insecticides).**

PHYTOSANITARY TREATMENT PROGRAM 2026/2027 SEASON

First Window

Two (2) treatments **T1** and **T2** (or three : **T1**, **T2** and **T3**) are made with alternative products to pyrethroids.

Country	Product (Trade name)	Formulation (Active ingredient)	Rate of use (L or g/Ha)
Bénin	JACOBIA 350 EC	Pyriproxifène 150 g/l + Nitempyrame 150 g/l	0,5
	FLOMID	Flonicamide 500 g/kg	100 g
	JACOBIA SUPER 350 EC	Pyriproxifène 150 g/l + Nitempyrame 150 g/l + Indoxacarb 50g/l	0,5
	THALIS 112 EC	Emamectine benzoate 48 g/l - acetamipride 64 g/l	0,25
	CORAGEN 200 SC	Chlorantraniliprole 200 g/l	0,1
	SAFARI G	Indoxacarb- Acétamipride 120-64 g/l	0,25
	BELT EXPERT 480 SC	Flubendiamide (240) thiachlopride (240)	0,1
	DRONE S 180 OD	Chlorantraniliprole (100) Spirotétramate (80)	0,125
	DRONE 200 OD	Chlorantraniliprole (200)	0,25
Burkina Faso	EMACOT PRO 300 WG	Emamectine benzoate 300g/kg	0,04
	JAZZFIRE 500 WG ; FLOCA 50 WG	Flonicamide 500g/kg	0,1
	VELEKTIN 365 WG	Emamectine benzoate 15 g/kg + profenofos 350 g/kg	0,7
	AVAUNT 150 EC	Indoxacarbe 150 g/l	0,17
Cameroun	THIAN 175 O -Teq	Flubendiamide + spirotétramate	0,2
	SPANVIC PLUS	Imidaclopride + Spinosad	0,4
	SILVALIS	Plinazoline	0,15
Côte d'Ivoire	FLOMID MAX 350 SC	Flonicamide 250 g/kg + Cyhalodiamide 100 g/kg	0,2
	JACOBIA 350 EC NOUVEAU	Pyriproxifène 150 g/l + Nitempyram 150 g/l + indoxacarde 50 g/l	0,5
	LEBON 184 WP	Emamectine benzoate 56 g/l + Acétamipride 128 g/l	0,25
	DANGORO PRO 168 EC ; FULL PRO 168 EC ; DANGORO PRO 168 DC	Emamectine benzoate 48 g/l + Acétamipride 120 g/l	0,25
	LOKI 420 WG	Emamectine benzoate 120 g/kg + Acétamipride 300 g/kg	0,25
	ZIDAN PRO 430 WG	Emamectine Benzoate : 80 g/kg Flonicamide : 350 g/kg	150 g
	THALIS PRO 168 EC	Emamectine 48 g/l + Acétamipride 120 g/l	0,25
	DENVER 666 WG	Chlorantraniliprole: 250 g/kg + Flonicamide : 416 g/kg	120 g
Mali	OGOUN 200SC ; CORANGEN 20 SC	Chlorantraniliprole 200g/l	0,1
	EMACOT 19 EC	Emamectine benzoate	0,5
	TIHAN 175 Q-TEQ ; MOVENTO TOTAL	Spirotetramate 75g/l + Flubendiamide 100g/l	0,2
	INDOX	Indoxacarb 150 g/l	0,17
	NOMOLTR 150 SC	Teflubenzuron	0,171
	COBRA 120 EC	Spinotoram 64 g/l + Acétamiprid 56g/l	0,25
	ULALA 500 WG	Flonicamide 500 g/kg	100 g
	FLYER PLUS 275 EC	Dinotéfuran 220 g/l + Bifenthrine 55 g/l	0,5
	JACOBIA ORDINAIRE 350 EC	Pyridine fonctionnelle 150 g/diamine fonctionnelle 200g/l	0,5
Sénégal	SILVALIS 200 SC	Plinazoline 200g/l	0,15
Tchad	THALIS 56 EC	Emamectine Benzoate 24g/l + Acétamipride 32g/l	0,5
	POLARIS 112 EC	Emamectine Benzoate 48 g/l + Acétamipride 64 g/l	0,25
	JAZZFIRE 500WG	Flonicamide 500 g/kg	100 g
Togo	TIHAN 175 OTEQ	Spirotetramate 75g/l + Flubendiamide 100g/l	0,2
	BELT EXPERT 480	Flubendiamide 240g/l + Tchiaclopride 240g/litre	0,1
	STINIPROL FORTE 350 SC	Chlorantraniliprole 100 g/L + Flonicamide 250g/l	0,2
	JAZZFIRE 500 WG ; NICOMEDE 500 WG	Flonicamide 500g/kg	100 g
	STINIPROLE 200 SC	Chlorantraniliprole 200g/l	0,1

Second window

Two treatments, **T3** and **T4** (or three **T4**, **T5** and **T6**) carried out following the 1st window.

Country	Product (Trade name)	Formulation (Active ingredient)	Rate of use (L or g/Ha)
Bénin	BELLA FTE 424 EC	Deltaméthrine-chlorpyriphos-éthyl 24-400g/l	0,5
	JACOBIA SUPER 350 EC	Pyriproxifène 150 g/l + Nitempyrame 150 g/l+ Indoxacarb 50g/l	0,5
	PYRO FTE 672 EC	Cyperméthrine-chlorpyriphos-éthyl 72-600 g/l	0,5
Burkina Faso	CORAGEN 20 SC	Chlorantraniliprole 200 g/l	0,1
	FLYER PLUS 275 EC	Dinotefuran 220 g/l + bifenthrine 55 g/l	0,5
	VELEKTIN 365 WG	Emamectine benzoate 15 g/kg + profenofos 350 g/kg	0,7
	AVAUNT 150 EC	Indoxacarbe 150 g/l	0,17
	DENVER666 WG ; DEVIL 666 WG	Chlorantraniliprole 250 g/kg + Flonicamide 416 g/kg	120 g
	LAUDENTO 200 SC	Plinazoline 200 g/l	0,15
Cameroun	CORAGEN + FLONIROY	Rynaxypyr + Flonicamide	150ml+100g
	IMPERATOR + OPTIMAL	Cyhalodiamide +Acétamipride	150ml+100g
	RADIAN + FLONIROY	Spinétoram + Flonicamide	120ml+100g
Côte d'Ivoire	TIHAN 175 OTEQ	Flubendiamid 100 g/l + Spirotetramat 75 g/l	0,2
	DENVER 666 WG ; ZIDAN STARR 666 WG	Flonicamide 416 g/Kg + Chlorantraniliprole 250 g/Kg	120 g
	ELEVEST 265 SC	Chlorantranilipole 106 g/l + Bifenthrine 159 g/	0,2
	INDOXAN DUO 220 EC	Indoxacarbe 120 g/l + Acetamipride 100 g/l	0,25
	JACOBIA 350 EC	Pyriproxifène 150 g/l + Nitempyram 150 g/l + indoxacarde 50 g/l	0,5
	GRACIA 10 EC	Fluxametadine 100 g/l	0,3
	ANGEL 320 SC ; STINGER EXTRA 320 SC ; BOOSTER FORT 320 SC	Flonicamide 200 g/l + Indoxacarbe 120 g/l	0,25
	KORONI FLA 360 SC	Flonicamide 200 g/l + Indoxacarbe 160 g/l	0,25
	FLOMID MAX 350 WG	Flonicamide 250 g/l + Cyhalodiamide 100 g/l	0,2
	JACOBIA 350 SC Nouveau	Pyriproxifène 150 g/l + Nytempyram 150 g/l + Indoxacarbe 50 g/l	0,5
	TSUNAMIDE 280 OD	Flonicamide 200 g/l + Flubendiamide 80 g/l	0,25
	NICKEL FORT 280 EC	Dinotéfurane 220 g/l + Indoxacarbe 60 g/l	0,5
	SILVALIS 200 SC	Plinazolin	0,15
Mali	ATTAKAN	Cyperméthrine 144g/l+Imidachlopride 200g/l	0,25
	IMUNIT 150 SC	Alphacyperméthrine 75g/l +Téflubenzuron 75g/l	0,2
	THALIS 112 EC	Emamectine benzoate 48g/l+acétamipride 64g/l	0,25
	PYRIMECTINE ; EMAPYR 80 EC	Emamectine benzoate /Pyriproxifène 20/60	0,5
	TAGANA 104 EC ; TOUMOFLA 104 EC ; EMIR FORT 104 EC	Cyperméthrine 144 g/Acétamipride 72/32	0,5
	GRACIA 10 EC	Fluxametamide 100 g/l	0,3
	JACOBIA NOUVEAU	Pyridine fonctionnelle 150 g/diamine fonctionnelle 150+indoxacarbe 50 g/l	0,5
Sénégal	JACOBIA 350 EC SUPER	Nitenpyrame150 g/l + Pyriproxifene 150 g/l + Indoxacarbe 50 g/l	0,5
Tchad	OGOUN 200 SC	Chlorantraniliprole 200g/l	0,15
	JAZZFIRE 500WG	Flonicamide 500 g/kg	100 g
Togo	ALFAZOL 96 EC 0,5l/ha	Alphacyperméthrine 36g/litre + Etoxazole 60g/litre	0,5
	FLYER PLUS 275 EC	Bifenthrine 55g/l + Dinotéfurane 220g/l	0,5
	JAZZFIRE 500 WG ; NICOMEDE 500	Flonicamide 500g/kg	100 g

Third window

Two treatments, **T5** and **T6** (or three **T7**, **T8** and **T9**) carried out following the 2nd window

Country	Product (Trade name)	Formulation (Active ingredient)	Rate of use (L or g/Ha)
Bénin	THALIS 112 EC	Emamectine benzoate 48 g/l - acetamipride 64 g/l	0,25
	SAFARI GOLD 184 EC	Indoxacarb (120)-acetamipride (64)	0,25
	CORAGEN 200 SC	Chlorantraniliprole 200 g/l	0,1
	DRONE S 180 OD	Chlorantraniliprole (100) Spirotétramate (80)	0,125
	DRONE 200 OD	Chlorantraniliprole (200)	0,25
	JACOBIA SUPER 350 EC	Nitenpyrame 150 g/l + Pyriproxyfene 150 g/l + Indoxacarbe 50 g/l	0,5
	COTONIX 328 EC	Deltaméthrine 12 g/l - Chlorpyriphos-éthyl 300 g/l - acétamipride 16 g/l	1
Burkina Faso	LAUDENTO 200 SC	Isocycloseram (Plinazoline) 200 g/l	0,15
	ANGEL 320 SC	Indoxacarbe 120 g/l + flonicamide 200 g/l	0,25
	DENVER 666 WG	Chlorantraniliprole 250 g/kg + flonicamide 416 g/kg	120 g
	CAPT SUPER 104 EC	Cyperméthrine 72g/l+ Acétamipride 32 g/l)	0,5
	RAZZIA 208 EC	Cyperméthrine 144 g/l+ Acétamipride 64 g/l	0,25
	SAUVEUR 62 EC	Lambdacyhalothrine 30 g/l + acétamipride 32 g/l	0,5
Cameroun	MORAN30DF + OPTIMAL	Indoxacarbe + Acétamipride	84 g+50 g
	DINEL	Emamectine + Téflubenzuron	200 g
	EMACOT PRO	Emamectine	60 g
	AZERTY	Emamectine	0,12
Côte d'Ivoire	DENVER 666 WG ; ZIDAN STAR 666 WG ; TONGA 666 WG	Chlorantraniliprole 250 g/Kg + Flonicamide 416 g/Kg	120 g
	INDOXANDUO 220 EC ; JAGUAR SUPER 220 EC	Indoxacarbe 120 g/l + Acetamipride 100 g/l	0,25
	BELT EXPERT 480 SC	Flubendiamide 240 g/l + Thiaclopride 240 g/l	0,1
	IBIS A 68 EC	Alphacyperméthrine 36 g/l + Acétamipride 32 g/l	0,5
	FARAFINA 104 EC	Cyperméthrine 72 g/l Acétamipride 32 g/l	0,5
Mali	CYPERPRO 720 EC	Cypermétrine 144g/l+Imidaclopride 200g/l	0,250
	GRACIA 10 EC	Fluxametamide 100 g/l	0,3
	JACOBIA NOUVEAU	Pyridine fonctionnelle 150 g/diamine fonctionnelle 150+indoxacarbe 50 g/l	0,5
Sénégal	DEVIL 666 WG	Flonicamide 416 g/kg + Chlorantraniliprole 250 g/kg	120 g
Tchad	EMIR FORT 104 EC	Cyperméthrine 72 g/l et Acétamipride 32 g/l EC	0,5
	JAZZFIRE 500WG	Flonicamide 500 g/kg	100 g
Togo	ATTAKAN C 344 SE	Cyperméthrine 144g/l + imidaclopride 200g/	0,25
	DELTAPYR 84 EC	Deltaméthrine 24g /l + pyriproxyfène 60g/l	0,5
	JAZZFIRE 500 WG ; NICOMEDE 500 WG	Flonicamide 500g/kg	100 g

REMINDER OF EFFECTIVE ACTIVE INGREDIENTS RECOMMENDED BY PR-PICA AGAINST JASSIDS (*A. biguttula*)

Active ingredients	g or ml p.c/ha	Effectiveness/pest						
		Jassid	Puc	Bem	Helico	Earias	Thau	Pect
Plinazoline 200 g/l (SC)	150	Effective	Not effective	Not effective	Effective	Effective	Moderately Effective	Moderately Effective
Alphacypermethrine 45 g/l + Dinotéfurane 200 g/l (SC)	400	Effective	Moderately Effective	Moderately Effective	Effective	Effective	Effective	Effective
Chlorfenapyr 250 g/l + Dinotéfurane 150 g/l (SC)	500	Effective	Effective	Moderately Effective	Effective	Effective	Not effective	Not effective
Nitenpyrame 200 g/l + Pyriproxyfène 150 g/l (EC)	500	Effective	Effective	Effective	Not effective	Not effective		
Nitenpyrame 150 g/l + Pyriproxyfène 150 g/l + Indoxacarbe 50 g/l (EC)	500	Effective	Effective	Effective	Effective	Effective		
Dinotéfurane 220 g/l + Bifenthrine 55 g/l (EC)	500	Effective	Effective	Effective	Moderately Effective	Effective		
Dinotéfurane 10% + Tolfenpyrad 15% (SC)	750	Effective	Effective	Effective	Moderately Effective			
Fluxamétamide 100 g/l (EC)	300	Effective	Effective	Moderately Effective	Effective	Effective		
Flonicamide 416 g/kg + Chlorantraniliprole 250 g/kg (WG)	120	Effective	Effective	Effective	Effective	Effective		
Flonicamide 200 g/l + Chlorantraniliprole 100 g/l (OD)	250	Effective	Effective	Effective	Effective	Effective		
Flonicamide 200 g/l + Cyhalodiamide 80 g/l (SC)	250	Effective	Effective	Moderately Effective	Effective	Effective		
Flonicamide 250 g/l + Cyantraniliprole 250 g/l (SC)	200	Effective	Effective	Effective	Effective	Effective		
Flonicamide 200 g/l + Flubendiamide 80 g/l (OD)	250	Effective			Effective	Effective		
Flonicamide 100 g/l + Indoxacarbe 60 g/l (SC)	500	Effective			Effective	Effective		
Flonicamide 200 g/l + Indoxacarbe 120 g/l (SC)	250	Effective	Effective	Effective	Effective	Effective	Moderately Effective	
Flonicamide 200 g/l + Pyriproxyfène 200 g/l (EC)	250	Effective	Effective	Effective	Not effective	Not effective	Not effective	Not effective
Nitenpyrame 150 g/l + Emamectine 24 g/l (OD)	500	Effective	Effective	Effective				
Flonicamide 500 g/kg (WG)	100	Effective	Effective	Not effective	Not effective	Not effective		
Flonicamide 500 g/l (OD)	100	Effective	Effective	Not effective	Not effective	Not effective		
Flonicamide 200 g/kg + Emamectine 60 g/kg (WG)	250	Effective	Moderately Effective		Effective	Effective		
Flonicamide 200 g/l + Emamectine 60 g/l (SC)	250	Effective	Effective		Effective	Effective		
Flonicamide 200 g/l + Emamectine 48 g/l (SC)	250	Effective	Effective		Effective			



Effective



Moderately Effective



Not effective



not available